



An Introduction to
Plate Tectonics

Prof. B. N. Upreti

Tribhuvan University
Arba Minvh University
University of Zambia


PLATE TECTONICS

CONTINENTAL DRIFT




The suggestion that there might have been lateral displacements of the continental masses on a gigantic scale is generally ascribed to ***F.B. Taylor*** in America (1908) and to ***Alfred Wegener*** in Germany (1910). For several years they developed their ideas independently.

However, the germ of the idea can be traced back to 1620, when ***Francis Bacon*** pointed out the parallelism of the opposing shores of the Atlantic.

Nearly two centuries later ***Antonio Snider*** united the continents in much the same way as Wegener did. He published a book in 1858 with two maps showing the reassembling of the continents.



A) Maps made by Antonio Snider-Pelligrini in 1858.



(B) Maps made by Alfred Wegener in 1915.

FIGURE 17.1 Continental drift was illustrated as early as 1858 by Antonio Snider-Pelligrini when he published these maps (A) in his book *Creation and Its Mysteries Revealed*. The idea seemed too far-fetched to the public and the scientific communities of the time and was forgotten, not to be revived for 50 years. Wegener published his series of maps (B) in 1915. His evidence, most of which was quite valid, was drawn from all of the sciences. Wegener called the original land mass Pangaea (“all lands”) and believed that the continents somehow plowed through the oceanic crust as they drifted.

Taylor (1910): He pictured the original Laurasia as being a continuous sheet of sial and supposed it to have spread outwards towards the equator, more or less radially from the polar region, much as a continental ice sheet would do. Wherever the resistance was least, the crust flowed out in lobes, raising up mountainous loops and arcs in front.

Evidences for continental drift

Wegener (1915)

- Paleontological evidence**
- Evidence from structure and rock types**
- Evidence from glaciation**
- Evidence from other paleoclimatic records**



Photo by Upreti taken from the exhibit of Natural History Museum, London)



■ **FIGURE 7.4** Bullard's fit of South America and Africa. The outlines correspond to the 914-m bathymetric contour. Light red shaded areas represent gaps and gray shaded areas represent overlaps in the fit.




Figure 19.3 This shows the best fit of South America and Africa along the continental slope at a depth of 500 fathoms (about 900 meters). The areas where continental blocks overlap appear in brown. (After A. G. Smith, "Continental Drift." In *Understanding the Earth*, edited by I. G. Gass. Courtesy of Artemis Press)



Alfred Wegener

Died at the age of 50



FIGURE 3.1 Alfred Wegener, the German meteorologist who proposed a comprehensive model of continental drift and presented geologic evidence in support of the idea.




FIGURE 3.2 Wegener's image of Pangaea and its subsequent breakup and dispersal.




PERMIAN
225 million years ago




TRIASSIC
200 million years ago




JURASSIC
135 million years ago




CRETACEOUS
65 million years ago




PRESENT DAY




200 million years ago



120 million years ago




56 million years ago



Present

Paleontological evidence






FIGURE 17.2 Paleontologic evidence of continental drift can be appreciated by considering the distribution of some fossil plants and animals found in South America, Africa, Madagascar, India, Antarctica, and Australia. *Mesosaurus*, a Permian freshwater reptile, is found in both Brazil and South Africa. *Glossopteris*, a fossil fern, is found on all of the southern continents in the zone shown on the map. *Lystrosaurus*, a Triassic land reptile, is found in South Africa, South America, India, and Antarctica. *Cynognathus*, an older Triassic reptile, is found in Argentina and South Africa. (Modified from L. Motz)




Figure 19.4 Fossils of *Mesosaurus* have been found on both sides of the South Atlantic and nowhere else in the world. Fossil remains of this and other organisms on the continents of Africa and South America appear to link these landmasses during the late Paleozoic and early Mesozoic eras.






Figure 19.5 These sketches by John Holden illustrate various explanations for the occurrence of similar species on landmasses that are presently separated by vast oceans. (Reprinted with permission of John Holden)

Evidence from paleoclimatic records

FIGURE 3.4 Map of Pangaea, showing the distribution of coal deposits and reefs (indicating tropical environments), and sand-dune deposits and salt deposits (indicating subtropical environments). Note how deposits now on different continents align in distinct belts.








FIGURE 17.6 Paleoclimatic evidence for continental drift includes deposits of coal, desert sandstone, rock salt, wind-blown sand, gypsum, and glacial deposits. Each indicates a specific climatic condition at the time of its formation. The distribution of these deposits is best explained if we assume that the continents were grouped together at the end of the Paleozoic Era, as shown in this diagram. (After American Association of Petroleum Geologists)

Evidence from glaciation



A



B

Figure 17.1 Pangaea The continents attained their present shapes when Pangaea broke apart 200 million years ago. A. The shape of Pangaea, determined by fitting together pieces of continental crust along a contour line 2000 m below sea level. (This is the line halfway down the continental slope, along which continental crust meets oceanic crust.) In a few places in this drawing, some overlap (black) occurs; elsewhere, small gaps (red) are found. These are places where post break-up

events have modified the shapes of the continental margins. The white area is the region affected by continental glaciation 300 million years ago. B. Present continents and the 2000-m contour below sea level. The white areas are where evidence of the old ice sheets exists. Arrows show directions of movement of the former ice. The dashed line joining the glaciated regions indicates how large the ice sheet would have to have been if the continents were in their present positions at the time of glaciation.

Evidence from structure and rock types










FIGURE 17.4 When reconstructed, the continents fit together like a jigsaw puzzle or pieces of a torn newspaper. Not only do the outlines of the torn pieces fit together, but the printing on them (analogous to the ages and structural features of the continents) also matches across their edges.

FIGURE 3.6 (a) Distinctive areas of rock units on South America link with those on Africa, as if they were once connected and later broke apart. (b) If the continents are returned to their positions in Pangaea by closing the Atlantic, mountain belts of the Appalachians lie adjacent to similar-age mountain belts in Greenland, Great Britain, Scandinavia, and Africa.





■ **FIGURE 7.11** The opening of the Atlantic Ocean basin. (a) The modern Atlantic Ocean and surrounding continents. (b) A continental reconstruction of 130 million years ago. Compare with the configuration of Pangaea at 255 million years ago shown in Figure 7.10. Africa and South America have moved east relative to North America. The South Atlantic has begun to open although opening of the Central Atlantic is well under way, and the Paraná and Etendeka flood basalts have already erupted. The North Atlantic has not yet started to open. (c) The Atlantic as it looked 80 million years ago. The ocean basin separating Africa, South America, and North America has developed and widened. The rift between northern Europe and North America (Greenland) has yet to develop. The position for the rifting of 60 million years ago is indicated by the future location of the North Atlantic flood basalt province. (d) By 36 million years ago, the North Atlantic has opened and separated the North Atlantic flood basalt province.



(a)

(b)

(c)

PLATE TECTONICS

The period between 1950 and 1960 was very important in bringing the revolution in earthscience field

The study of the *geology of ocean floor* and the *paleomagnetic studies* of rocks from continents and oceans during the above decade revived the theory of continental drift.

Two studies in paleomagnetism were very important:

- 1. Apparent Polar Wandering**
- 2. Patterns of Magnetic Reversals on the Sea Floor**



H.H. Hess

The man who gave the theory of sea floor spreading

SEA-FLOOR SPREADING

In 1960 **H.H. Hess** from Princeton University (USA) proposed a theory of *sea-floor spreading* which took into account the new data from echo soundings and also suggested a possible mechanism for continental drift.

Rock magnetism: The study of rock magnetism developed during the 1950s with the perfection of new, highly sensitive magnetometers.


In 1963, **Fred Vine** (graduate student) and **D.H. Matthews** saw a way to test the idea of Sea-floor spreading put forth by Hess. If sea-floor spreading has occurred, they suggested, it would be recorded in the magnetism of the basalts in the oceanic crust. Investigations proved this theory.

Evidence from the ocean floor





East Pacific rise









FIGURE 3.21 (a) In a mid-ocean ridge, heat from the mantle flows up through the crust; heat flow decreases away from the ridge axis. (b) The graph shows heat flow as a function of distance from the ridge axis. Heat flow is measured in heat-flow units, or HFU ($1 \text{ HFU} = 10^{-6} \text{ calories/cm/s}$).




(a)





Evidence from sediments on the ocean floor

Glomar Challenger started drilling on the ocean bottom. Youngest sediments rest on the basalt of the ocean floor near the oceanic ridge, away from the ridge, the sediments that directly lie above the basalt becomes progressively older.






FIGURE 4.10 This map of the world shows the age of the sea floor. Note how the sea floor grows older with increasing distance from the ridge axis. (Ma = million years ago.)

Trenches





Figure 19.23 Distribution of the world's oceanic trenches, ridge system, fracture zones, and transform faults. Where transform faults offset ridge segments, they permit the ridge to change direction (curve) as can be seen in the Atlantic Ocean.



Central
America
trench


Puerto-Rico
trench

Mid-Atlantic
Ridge

Peru-Chile
trench

South Sandwich
trench

Lithosphere-Asthenosphere Interaction






FIGURE 18.21 The low velocity zone is revealed by a drop in the velocities of both P waves (shown here) and S waves. This marks a zone of low strength in the upper mantle between about 100 and 250 km below the surface. The low-velocity zone is contained in the asthenosphere and marks part of the mantle that is very near its melting point and may be a zone of partial melting.





FIGURE 18.22 Discontinuities in seismic wave velocities may correspond to phase changes. The blue line shows how P wave velocities change with depth. The other lines show velocities for various minerals. The uppermost mantle is dominated by olivine. Below 400 km a velocity increase implies olivine is replaced by spinel. At greater depths, spinel is probably replaced by magnesium-perovskite. Each change increases the density of the mantle.



■ **FIGURE 2.25** (a) A familiar example of convection, in which soup is heated in a pot. Hot soup at the bottom of the pot above the flame expands and rises because of its buoyancy. It cools upon reaching the surface and sinks, completing the cycle. Within Earth, the heat is mainly supplied by radioactivity. (b) When the material through which heat flows is strong, like the base of a pot, the heat is transported by conduction. Conduction is the mechanism by which heat is transported across the lithosphere—the strong outer shell of Earth.





FIGURE 2.10 Convection in the mantle can be compared to convection in a pot of soup. Heat from below causes the material to expand and thus become less dense. The warm material rises by convection and spreads laterally. It then cools, and thus becomes denser, and sinks. It is reheated as it descends, and the cycle is repeated.




Figure 1-14 Convection cells and plate motion. Heat within the Earth's mantle results in rising ("convecting") currents of warm mantle material, which drag the lighter lithospheric plates along with them as they flow beneath the Earth's surface. As rising mantle material spreads beneath the lithospheric plates, it cools, becomes denser, and sinks back to the deeper interior, where it is reheated to rise again. Such a cycle, known as a convection cell, may be the principal driving mechanism of plate tectonics.






Figure 19.12 Seafloor spreading. Harry Hess proposed that upwelling of mantle material along the mid-ocean ridge system created new seafloor. The convective motion of mantle material carries the seafloor in a conveyor-belt fashion to the deep-ocean trenches, where the seafloor descends into the mantle.




Shallow
convection-cell
hypothesis

(a)



Deep
convection-cell
hypothesis


(b)




Two-tiered
convection-cell
hypothesis

(c)


Figure 11-21 Hypotheses proposed to explain the configuration of the Earth's convection cells. (a) The cells may be shallow, confined to the asthenosphere. **(b)** The cells may extend through the entire mantle to the outer core. **(c)** Some hypotheses propose two sets of convection cells that meet at a depth of 670 kilometers (420 miles), the zone within the mantle where seismic waves are known to accelerate in response to changes in the chemistry and structure of mantle materials.



A.



B.



C.

Figure 19.32 Proposed models of the driving force for plate tectonics. **A.** Large convection cells in the mantle may carry the lithosphere in a conveyor-belt fashion. **B.** Slab-pull results because the subducting slab is more dense than the underlying material. Slab-push is a form of gravity sliding caused by the elevated position of lithosphere at a ridge crest. **C.** The hot plume model suggests that all upward convection is confined to a few narrow plumes, while the downward limbs of these convection cells are the cold, dense subducting oceanic plates.




FIGURE 18.23 Earth's thermal structure and convection can be modeled using computers to complement the observations of seismic tomography. In one model, subducted slabs pass without pausing through the phase boundary at 660 km. In another model, the phase boundary is a temporary barrier that is broken down when enough subducted material accumulates and then flushes rapidly through the lower mantle. The lower mantle may convect by generating thin plumes that rise off of the core-mantle boundary. Some of the plumes may be triggered by the sinking of the dense overlying mantle.




Figure 3.26 Mantle plumes play a key role in some models of whole-mantle convection. The plumes may rise to the surface as isolated hot spots or they may combine to initiate the crustal rifting that results in the formation of lithospheric plates.



Figure 2.9 Spreading Center Splitting a Continent The African Rift Valley, which extends from the Red Sea in the north to Malawi in the south, is a gigantic rent in the Earth's surface marking the place where a spreading center is splitting Africa into two pieces. This photo is of a portion of the Rift Valley in central Kenya. To the east (right-hand side) is a plateau bounded by a jagged fracture. The two hills (rear and left-hand edge) are volcanoes.






Figure 2.16 Midocean Ridges on Land Long, deep fractures that split Iceland mark the center of a midocean ridge. Iceland is on the Mid-Atlantic Ridge and is one of the few places in the world where the midocean ridge can be seen above sea level.





X Shallow earthquakes
(tension and normal faulting at divergent boundaries; strike-slip at transform faults)


PLATE 12

Earthquake sources associated with two types of plate boundaries: divergent boundaries of ocean ridges and transform faults. [From Press and Siever, *Understanding Earth*, Second ed., New York: W. H. Freeman and Co., 1998.]













FIGURE 2.8 The tectonic system is powered by Earth's internal heat. The asthenosphere is more plastic than either the overlying lithosphere or the underlying lower mantle. Above the plastic asthenosphere, relatively cool and rigid lithospheric plates split and move apart as single mechanical units (along the ocean ridges). As this happens, molten rock from the asthenosphere wells up to fill the void between the lithospheric plates and thus creates new lithosphere. Very slow convection occurs in the mantle. Some plates contain blocks of thick, lower-density continental crust, which cannot sink into the denser mantle. As a result, where a plate carrying continental crust collides with another plate, the continental margins are deformed into mountain ranges. Plate margins are the most active areas on Earth—the sites of the most intense volcanism, seismic activity, and crustal deformation. Locally, convection in the deep mantle creates rising mantle plumes. (After P. J. Wyllie)

Atlantic Ocean

Mid-Atlantic
Ridge

South
America


Africa



Eurasian Plate

Indian Plate

Subducting plate






PLATE 13

Shallow- and deep-focus earthquakes along a convergent slab plunging underneath the upthrown mountain range. Magma is rising from the slab boundary at depth.

[From Press and Siever, *Understanding Earth*, Second ed., New York: W. H. Freeman and Co., 1998.]




Figure 2.15 Three Plates Simplified diagram of a section across a portion of the Pacific Ocean margin of southern South America at about the latitude of Puerto Monte, Chile. Note that the side of the trench adjacent to the continent (to the right) is steeper than the oceanic side.








FIGURE 1.5 The outermost layers of the solid Earth, based on physical characteristics, are the asthenosphere and the lithosphere. The asthenosphere is hot, close to its melting point, and is capable of plastic flow. The lithosphere above it is cooler and rigid. It includes the uppermost part of the mantle and two types of crust: thin, dense oceanic crust and thick, buoyant continental crust.







Central Andes (higher and drier)



Southern Andes (lower and wetter)





FIGURE 2

Cross sections of the Andes at different latitudes. Comparison among the three sections shows varying slab dip: steep slab dip in (a) and (c) and shallow slab dip in (b). Note that there is no active volcanism where the slab dip is shallow (see Fig. 1). This kind of shallow subduction may cause compression in the upper plate (compare to Fig. 10.2), leading to thrusting and crustal thickening, which has already happened farther north (a), with a subsequent increase in slab dip to today's configuration.


30 million years ago




20 million years ago







10 million years ago



Present



EXPLANATION

-  Spreading center (divergent boundary)
-  Subduction zone (convergent boundary)
-  Transform fault, arrows show relative movement
SAFZ, San Andreas fault zone
-  Triple plate junction
M, Mendocino
R, Rivera





FIGURE 10.3 (a) Distribution of earthquakes in a subduction zone, which is related to zones of tension and compression in the subducted slab. At shallow depths, earthquake hypocenters are located near the top of the subducted lithosphere. At greater depths, earthquakes tend to occur within the subducted slab. Note that some earthquakes also occur in the upper plate, probably due to stresses generated by contact between the two plates. (b) Tomographic image of the subduction zone beneath Japan. A color-coded scale defines the percentage of deviation in P-wave velocity from an assumed average value. Blue colors indicate fast seismic velocities (velocities that are up to 6 percent faster than the assumed average), which occur in the slab because it is relatively cool and dense. Yellow and red colors indicate P-wave velocities that are slower than the assumed average, which occur particularly beneath volcanoes. This probably indicates partial melting of the mantle, which would slow the velocity of the seismic waves. Earthquake locations are marked by the open circles.


Rock magnetism

The study of rock magnetism developed during the 1950s with the perfection of new, highly sensitive magnetometers. Two studies in paleomagnetism were very important:

- 1. Apparent Polar Wandering**
- 2. Patterns of Magnetic Reversals on the Sea Floor**



A.





B.


Figure 19.11 Simplified apparent polar-wandering paths as established from North American and Eurasian paleomagnetic data. **A.** The more westerly path determined from North American data was caused by the westward movement of North America by about 24 degrees from Eurasia. **B.** The positions of the wandering paths when the landmasses are reassembled.

In 1963, **Fred Vine** (a graduate student) **and** **D.H. Matthews** (supervisor) saw a way to test the idea of Sea-floor spreading put forth by Hess. If sea-floor spreading has occurred, they suggested, it would be recorded in the magnetism of the basalts in the oceanic crust. Investigations proved this theory.

A. Magnetometer record showing symmetrical magnetic field across ridge



B. Research vessel towing magnetometer across ridge crest



C. Location map

Figure 19.15 The ocean floor as a magnetic tape recorder. **A.** Schematic representation of magnetic intensities recorded as a magnetometer is towed across a segment of the Mid-Atlantic Ridge. **B.** Notice the symmetrical stripes of low- and high-intensity magnetism that parallel the ridge crest. Vine and Matthews suggested that the stripes of high-intensity magnetism occur where normally magnetized oceanic basalts enhance the existing magnetic field. Conversely, the low-intensity stripes are regions where the crust is polarized in the reverse direction, which weakens the existing magnetic field.




FIGURE 7.7 Magnetic reversals recorded in rocks of the oceanic crust. (a) Location of a magnetic survey across the Reykjanes Ridge—part of the Mid-Atlantic Ridge—near Iceland. Dark-shaded bands represent crust with a normal polarity magnetic field; light blue-shaded areas are reversed. (b) Magnetic-field intensity variations along the survey line. High- and low-intensity fields correspond to normal and reversed polarity, respectively, as summarized in the strip at the bottom of the figure. Ages are determined by correlation with sequences of lavas on land (Fig. 7.6), which were dated isotopically using the potassium-argon (K-Ar) method.









Figure 19.16 As new basalt is added to the ocean floor at mid-ocean ridges, it is magnetized according to Earth's existing magnetic field. Hence, it behaves much like a tape recorder as it records each reversal of the planet's magnetic field.



A. Period of normal magnetism



B. Period of reverse magnetism



C. Period of normal magnetism




Figure 17.4 Oceanic Crust Records Magnetic Reversals Schematic diagram of oceanic crust. Lava extruded along an oceanic ridge forms new oceanic crust. As the lava cools, it becomes magnetized with the polarity of the Earth's magnetic field. Successive strips of oceanic crust have alternate normal (green) and reversed (brown) polarity. The ages of the reversals are the same as those in the magnetic polarity time scale shown in Figure 8.19.

PLATES

The outer rigid layer of earth - the ***lithosphere*** is divided into a mosaic of seven major plates and a number of smaller subplates.

Plate boundaries are the most significant structural elements of the planet Earth because they reflect the internal dynamics of the planet.

The new geography of ***tectonic plates*** is much different than the classical physical geography of our planet (distribution of continents and oceans) and more important in Geology.





FIGURE 4.3 The major plates making up the lithosphere. Note that some plates are all ocean floor, while some contain both continents and oceans. Thus, some plate boundaries lie along continental margins (coasts), while others do not. For example, the eastern border of South America is not a plate boundary, but the western edge is.









FIGURE 4.5 The locations of most earthquakes fall in distinct bands. These earthquake belts define the positions of the plate boundaries.





A.

Figure 12.3 Continental Plateaus Along Convergent Plate Boundaries Color-coded elevation maps of (A) the Tibetan Plateau, formed along the collision between the Australian-Indian



B.

Plate and the Eurasian Plate; and (B) the Altiplano (“high face” in Spanish), formed along the subduction boundary between the Nazca Plate and the South American Plate.



PLATE 10


Seismicity of the Earth, 1964 to 1995 (magnitudes greater than 5.1), with tectonic plate boundaries superimposed. The large circles are major earthquakes with a magnitude over 6.4. This Pacific projection is centered on the Indo-Australia plate. [Courtesy of International Association of Seismology and the Earth's Interior.]

PLATE 11

Projection of the Atlantic hemisphere showing the America and Africa plates, and the mid-Atlantic seismically active ridge. The seismicity and plate boundaries are similar to those in Plate 10. [Courtesy of International Association of Seismology and the Earth's Interior.]








American Plate


Eurasian Plate




Indian-Australian plate

Pacific plate

Photo taken by Upreti. An
outcrop of the very sharp
plate boundary in New
Zealand




**500 million
years ago**



**420 million
years ago**



**320 million
years ago**



**225 million
years ago**

Figure 11-20 The origin of the supercontinent Pangaea.






Figure 19.31 Several views of the breakup of Pangaea over a period of 200 million years.



D. 50 Million Years Ago (Early Cenozoic)



E. Present




Figure 19.C The world as it may look 50 million years from now. (From "The Breakup of Pangaea," Robert S. Dietz and John C. Holden. Copyright 1970 by Scientific American, Inc. All rights reserved.)

Plate Boundaries

Three kinds of plate boundaries are recognized and define three fundamental kinds of deformation and geologic activity:

1. Divergent Plate Boundaries:

Zones of tension, where plates split and spread apart

2. Convergent Plate Boundaries:

Zones where plates collide and (*Subduction Zones*) one plate moves down into the mantle

3. Transform Fault Boundaries:

Zones of shearing, where plates slide past each other without diverging or converging








Figure 19.18 Schematic representation of plate boundaries showing the relative motion of plates. **A.** Divergent boundary. **B.** Convergent boundary. **C.** Transform fault boundary.

Type of Margin	Divergent	Convergent	Transform
Motion	Spreading	Subduction	Lateral sliding
Effect	Constructive (oceanic lithosphere created)	Destructive (oceanic lithosphere destroyed)	Conservative (lithosphere neither created or destroyed)
Topography	Ridge/Rift	Trench	No major effect
Volcanic activity?	Yes	Yes	No






A. Divergent boundary





B. Convergent boundary



C. Transform fault boundary

Figure 1.16 Mosaic of rigid plates that constitute Earth's outer shell. (After W. B. Hamilton, U.S. Geological Survey)







Figure 2.22 Volcanism and Hot Spots Most volcanism on Earth occurs underwater at the midocean ridges. Most volcanoes above sea level are associated with subduction zones. In addition, hot spot volcanism associated with rising mantle

plumes (yellow circles) has been identified at several locations. Some hot spots, such as under Iceland, are associated with plate boundaries. Other hot spots, such as Hawaii, lie within plate interiors.



Mediterranean Sea

Arabian Peninsula

Red Sea

Triple junction

Africa

Gulf of Aden

Indian Ocean

Lake Turkana


Mt. Kilimanjaro

East African Rift

Lake Victoria

Lake Tanganyika

Lake Malawi



Thank you